

IPBF FACT SHEET

KETAMINE ABUSE AND THE URINARY TRACT

WHAT IS KETAMINE

Ketamine is a dissociative anaesthetic developed in the mid 1960s and is used in human and veterinary medicine. In medical settings, administration of ketamine may be intravenous, subcutaneous, intramuscular, oral, topical intranasal or sublingual. It is used for inducing and maintaining anaesthesia, for analgesia in a variety of pain settings, and as a rapid effect antidepressant. It is perhaps best known for its use in treating injured soldiers during the Vietnam War. Reported side effects in some patients have included hallucinations, nightmares, sedation, dizziness, blurred vision, agitation and nausea/vomiting. It may increase blood pressure and heart rate.

RECREATIONAL ABUSE

Nonmedical (illicit) use was first documented in the late sixties/early seventies when it was emerging as a club drug. By the nineties, ketamine abuse was rapidly increasing in East Asia, particularly in the dance culture of Hong Kong, while today it is being used recreationally worldwide due to its relative low cost compared to similar drugs. As a party drug, it is being used by young people, including very young teenagers. It may be snorted, injected or taken orally in pill form, sometimes masquerading as XTC pills. It may also be mixed with other drugs and of course accompanied by a lot of alcohol. Since ketamine is tasteless and odourless, it can be placed in drinks without the intended victim suspecting and as such is used as a “date rape drug”. The numerous street names for ketamine include K, Special K, Vitamin K, Super Acid, Super C, Bump, Cat Valium, Green K, Honey Oil, Special La Coke, Ket, Kitty, Kit Kat, New Ecstasy, Purple and Jet. A mixture of ketamine and cocaine is called Calvin Klein or CK1. In Hong Kong, the street name is Kai-Jai.

EFFECTS OF KETAMINE ABUSE

Effects associated with recreational ketamine abuse include a hallucinogenic-like effect, a pleasant dream-like state, sense of floating and being separated from your body. However, some ketamine “bad trips” can cause a frightening sensation of total dissociation compared to a near-death experience and known as the “K-hole”, with the sensation of being in a dark tunnel. In addition to the side effects mentioned above, ketamine abuse may also cause amnesia, flashbacks, memory impairment, anxiety, impaired motor function, respiratory and gastrointestinal disorders. In the past decade, there has been increasing realisation that ketamine abuse can lead to pain and damage to the urinary tract, including interstitial cystitis-like inflammatory bladder changes and symptoms, with an increased urge to urinate, blood in the urine, incontinence and pain on urination.

CLINICAL PRESENTATION

Patients may present with IC-like symptoms. Young people especially should therefore be questioned regarding the possibility of recreational ketamine abuse. Recent studies in different parts of the world have reported chronic inflammation, often severe lesions without malignancy, denuded epithelium, thickened bladder wall, contracted bladder, reduced bladder capacity, ureteric wall thickening, narrowing and strictures, swollen kidneys. The longer the drug abuse, the worse the risk of extensive damage.

TREATMENT

Cessation of ketamine abuse has been shown to reduce urinary tract symptoms in many of these patients. However, the outcome of treatment depends on the severity of the disease process. It is still unknown whether severe damage to the urinary tract is fully reversible. Long-term abusers may have such extensive damage to the bladder that cystectomy may be the only option. Results of

surgery may, however, be compromised if ketamine abuse is resumed or continued. Ketamine is a very addictive drug and if a patient has been using it for several years, they may find it impossible to stop. Treatment may additionally be complicated by the fact that after several years of ketamine abuse, patients may have developed many other physical and mental problems including cognitive impairment and schizophrenia-like symptoms. Since the mechanism of ketamine-associated urinary tract symptoms is as yet unknown, treatment is aimed at alleviating the symptoms, similar to interstitial cystitis. Treatments that protect the bladder lining such as intravesical sodium hyaluronate or pentosan polysulfate have been used with some success.

PREVALENCE

Prevalence of urinary tract symptoms among ketamine abusers is uncertain, but studies have indicated that between a quarter and one third of ketamine abusers may be affected.

AWARENESS

It is vital to raise awareness and to warn teenagers and young adults of the risks. Furthermore, if ketamine abuse continues to escalate, the resultant urinary tract symptoms could place high demands on health resources at substantial cost in the future.

References and further information:

- Chen CH, Lee MH, Chen YC, Lin MF. Ketamine-snorting associated cystitis. *J Formos Med Assoc.* 2011 Dec;110(12):787-91.
- Chu PS, Kwok SC, Lam KM, Chu TY, Chan SW, Man CW, et al. 'Street ketamine'-associated bladder dysfunction: A report of ten cases. *Hong Kong Med J.* 2007;13:311_3.
- Chu PS, Ma WK, Wong SC, Chu RW, Cheng CH, Wong S, et al. The destruction of the lower urinary tract by ketamine abuse: A new syndrome? *BJU Int.* 2008;102:1616_22.
- Cottrell A, Warren K, Ayres R, Weinstock P, Kumar V, Gillatt D. The destruction of the lower urinary tract by ketamine abuse: A new syndrome? *BJU Int.* 2008;102:1178_9.
- Cottrell, A. M, et al. "Urinary tract disease associated with chronic ketamine use". *British Medical Journal.* 3 May 2008.
- Curran HV, Monaghan L. (May 2001). "In and out of the K-hole: a comparison of the acute and residual effects of ketamine in frequent and infrequent ketamine users". *Addiction* 96 (5): 749-760.
- Jansen, Karl (2001). *Ketamine: Dreams and Realities.* Multidisciplinary Association for Psychedelic Studies. p. 24.
- Lai Y, Wu S, Ni L, Chen Z, Li X, Yang S, Gui Y, Guan Z, Cai Z, Ye J. Ketamine-associated urinary tract dysfunction: an underrecognized clinical entity. *Urol Int.* 2012;89(1):93-6.
- Mak SK, Chan MT, Bower WF, Yip SK, Hou SS, Wu BB, Man CY. Lower urinary tract changes in young adults using ketamine. *J Urol.* 2011 Aug;186(2):610-4.
- Mason K, Cottrell AM, Corrigan AG, Gillatt DA, Mitchelmore AE. Ketamine-associated lower urinary tract destruction: a new radiological challenge. *Clin Radiol.* 2010 Oct;65(10):795-800.
- Morgan, Celia J. A.; Muetzelfeldt, Leslie; Curran, H. Valerie (2009). "Consequences of chronic ketamine self-administration upon neurocognitive function and psychological wellbeing: a 1-year longitudinal study". *Addiction* 105 (1): 121-33.
- Pal R, Balt S, Erowid E, Erowid F, Baggott MJ, Mendelson J, Galloway GP. Ketamine is associated with lower urinary tract signs and symptoms. *Drug Alcohol Depend.* 2013 Mar 6. pii: S0376-8716(13)00042-2. doi: 10.1016/j.drugalcdep.2013.02.005. [Epub ahead of print]
- Selby NM, Anderson J, Bungay P, Chesterton LJ, Kolhe NV. Obstructive nephropathy and kidney injury associated with ketamine abuse [case report]. *Nephrology Dialysis Transplantation Plus.* 2008;1:310_2.
- Shahani R, Streutker C, Dickson B, Stewart RJ. Ketamine associated ulcerative cystitis: A new clinical entity. *Urology.* 2007;69:810_2.
- Tsai TH, Cha TL, Lin CM, Tsao CW, Tang SH, Chuang FP, et al. Ketamine-associated bladder dysfunction. *Int J Urol.* 2009;16:826_9.
- Winstock AR, Mitcheson L, Gillatt DA, Cottrell AM. The prevalence and natural history of urinary symptoms among recreational ketamine users. *BJU Int.* 2012 Dec;110(11):1762-6.
- Yeung LY, Rudd JA, Lam WP, Mak YT, Yew DT. Mice are prone to kidney pathology after prolonged ketamine addiction. *Toxicol Lett.* 2009;191:275_8.
<http://www.thegoooddrugsguide.com/ketamine/faq.htm>
http://en.wikipedia.org/wiki/Recreational_use_of_ketamine
<http://en.wikipedia.org/wiki/Ketamine>

Author: Jane M. Meijlink, Rotterdam, The Netherlands

Contact and further information: info@painful-bladder.org

The International Painful Bladder Foundation does not engage in the practice of medicine. It is not a medical authority nor does it claim to have medical knowledge. The IPBF endeavours to ensure that all information it provides is correct and accurate, but does not accept any liability for errors or inaccuracies.